

Clinical Trial Protocol

Iranian Registry of Clinical Trials

10 Jun 2026

Comparing the effectiveness or ineffectiveness of Valsalva maneuver and EMLA cream on level of venous cannulation pain

Protocol summary

Summary

The present study aims to compare the effect of Valsalva Maneuver and EMLA cream on level of venous cannulation pain. It is a single blind experimental study. 160 patients candidate to general surgeries including ear, nose and throat, ophthalmology and urology bedridden in Department of Internal Medicine, Hazrat Rasul hospital, Javanrud, Kermanshah were selected and randomly divided into 3 experiments groups and a control group. The first experiment group made use of EMLA cream, 30 mg. After 40 minutes of applying the cream, venous cannulation was accomplished. Valsalva Maneuver was done on the second group through blowing in a plastic pipe connected to a Mercury barometer, and Valsalva Maneuver was accomplished for at least 20 seconds. Venous cannulation was accomplished immediately after that. In the third group, both EMLA cream and Valsalva Maneuver were conducted so that EMLA cream was applied and, after 40 minutes, Venous cannulation was accomplished immediately in conditions similar to the first group. Vaseline was applied to the control group, and Venous cannulation was conducted after 40 minutes. Then, barometer was used to check patients' blood pressure. Heart beat and the percentage of arterial oxygen saturation were checked using Pulse Oximeter. Pain rate was measured 5 minutes before Venous cannulation and immediately after that. Inclusion criteria: avoiding Systemic analgesics; lack of neurologic defects; lack of respiratory, cardiac, glaucoma diseases; no skin allergy or Atopic dermatitis. Exclusion criteria: Intravenous cannula failure in the first attempt.

General information

Acronym

IRCT registration information

IRCT registration number: **IRCT2015062814333N37**

Registration date: **2015-07-24, 1394/05/02**

Registration timing: **registered_while_recruiting**

Last update:

Update count: **0**

Registration date

2015-07-24, 1394/05/02

Registrant information

Name

Feizollah Foroughi

Name of organization / entity

kermanshah University of Medical Sciences

Country

Iran (Islamic Republic of)

Phone

+98 83 1821 4653

Email address

fforoughi@kums.ac.ir

Recruitment status

Recruitment complete

Funding source

Vice chancellor for research, Kermanshah University of Medical Sciences

Expected recruitment start date

2015-07-23, 1394/05/01

Expected recruitment end date

2015-09-23, 1394/07/01

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Comparing the effectiveness or ineffectiveness of Valsalva maneuver and EMLA cream on level of venous cannulation pain

Public title

A Comparative study of the effect of Valsalva maneuver and EMLA cream on level of venous cannulation pain

Purpose

Other

Inclusion/Exclusion criteria

Inclusion criteria: avoiding Systemic analgesics; lack of neurologic defects; lack of respiratory, cardiac, glaucoma diseases; no skin allergy or Atopic dermatitis. Exclusion criteria: Intravenous cannula failure in the first attempt.

Age

From **15 years** old to **60 years** old

Gender

Both

Phase

N/A

Groups that have been masked

No information

Sample size

Target sample size: **160**

Randomization (investigator's opinion)

Randomized

Randomization description**Blinding (investigator's opinion)**

Single blinded

Blinding description**Placebo**

Not used

Assignment

Parallel

Other design features

Randomized by sealed envelope

Secondary Ids

empty

Ethics committees**1****Ethics committee****Name of ethics committee**

Vice chancellor for research, Kermanshah University of Medical Sciences

Street address

Building No.2, Shahid Beheshti, Vice Chancellor for Research Affairs, Kermanshah University of Medical Sciences

City

Kermanshah

Postal code**Approval date**

2015-05-26, 1394/03/05

Ethics committee reference number

Kums.Rec.1394.21

Health conditions studied**1****Description of health condition studied**

level of venous cannulation pain

ICD-10 code

Y60.6

ICD-10 code description

During aspiration, puncture and other catheterization

Primary outcomes**1****Description**

Pain level

Timepoint

Immediately after catheter insertion

Method of measurement

Using NRS according to criterion score gained from 0 to 10

2**Description**

Average venous pressure

Timepoint

5 minutes before venous cannulation and immediately after that

Method of measurement

Using barometer

3**Description**

Heart beat

Timepoint

5 minutes before venous cannulation and immediately after that

Method of measurement

Using pulse oximeter

4**Description**

Arterial oxygen saturation

Timepoint

5 minutes before venous cannulation and immediately after that

Method of measurement

Using pulse oximeter

Secondary outcomes

empty

Intervention groups**1****Description**

The first group made use of EMLA cream, made by Astrazeneca,Eczacibasi Ltd, Turkey. Cream surface is covered by a piece of paper and a 2 cc syringe was used

to apply 1 cc, or 2.5 mlg, of the cream on dependent hand where venous cannulation was conducted at 2.5×2.5 cm. after 40 minutes, the dressing was removed from cannulation point and the place was disinfected using cotton drained in alcohol. Then, BP18 IVC was used to do venous cannulation while the patient lay back and his head is against the cannulation point. Also, tourniquet was placed on his forearm.

Category

Treatment - Drugs

2

Description

Valsalva maneuver in the second group was accomplished through blowing in a plastic pipe connected to a mercury barometer (EasyLife, China) and holding up mercury bar upper than 20 mmHg for at least 20 seconds. Immediately after that venous cannulation was done similar to the group using EMLA cream.

Category

Treatment - Drugs

3

Description

Both EMLA cream and Valsalva maneuver were applied to the third group. 40 minutes after applying EMLA cream and in similar conditions to the first group, venous cannulation was done immediately after Valsalva maneuver.

Category

Treatment - Drugs

4

Description

All the conditions to the group using EMLA cream were considered to the fourth group. The difference is that Vaseline (Kimiagar-Toos Ltd, Iran) is used in the fourth group instead of EMLA cream. After 40 minutes venous cannulation was accomplished.

Category

Placebo

Recruitment centers

1

Recruitment center

Name of recruitment center

Hazrat Rasool Hospital, Javanrod

Full name of responsible person

Mahdi Babaei

Street address

Hazrat Rasool Hospital, Street of Baradarn Amini,
Javanrod

City

Javanrod

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Vice chancellor for research, Kermanshah University of Medical Sciences

Full name of responsible person

Koroush Hamzehee

Street address

Building No.2, Shahid Beheshti, Vice Chancellor for Research Affairs, Kermanshah University of Medical Sciences

City

Kermanshah

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Vice chancellor for research, Kermanshah University of Medical Sciences

Proportion provided by this source

100

Public or private sector

empty

Domestic or foreign origin

empty

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

empty

Person responsible for general inquiries

Contact

Person responsible for scientific inquiries

Contact

Name of organization / entity

Kermanshah University of Medical Sciences

Full name of responsible person

Dr. Rostam Jalali

Position

Assistance professor of Nursing Training

Other areas of specialty/work

Street address

School of Nursing & Midwifery, First of Dolatabad
Boulevard, Eisar Square

City

Kermanshah

Postal code

Phone

+98 83 3827 9304

Fax

Email

KS_jalali@yahoo.com

Web page address	<i>empty</i>
Person responsible for updating data	Statistical Analysis Plan <i>empty</i>
Contact	Informed Consent Form <i>empty</i>
Sharing plan	Clinical Study Report <i>empty</i>
Deidentified Individual Participant Data Set (IPD)	Analytic Code <i>empty</i>
Study Protocol	Data Dictionary <i>empty</i>