

# Clinical Trial Protocol

## Iranian Registry of Clinical Trials

10 Jun 2026

### The effects of Mobilization with movement technique and knee osteoarthritis conventional physiotherapy on disability, pain and spatiotemporal and kinematic characteristics of gait in individuals with knee osteoarthritis

#### Protocol summary

##### Study aim

General aim Investigating the effect of 10 treatment sessions of mulligan technique along with common physiotherapy in the knee joint on disability, pain and kinematic and spatiotemporal characteristics of walking and electromyographic activity of muscles during gait, sit to stand and static balance in people with knee osteoarthritis

##### Design

Clinical trial with treatment and control group, double-blind, stratified randomization on 38 patients by RANDOM NUMBER GENERATOR

##### Settings and conduct

First, the participants will be evaluated in the biomechanics laboratory, the subject and the examiner will be blind to the grouping and the examiner will not be present in the treatment process, then the examinees will receive 10 treatment sessions in the hospitals and clinics affiliated to the Iran University of Medical Sciences, then again will be evaluated in the laboratory.

##### Participants/Inclusion and exclusion criteria

- Age 40 to 70 years -Diagnosis of knee tibiofemoral joint arthrosis based on American College of Rheumatology criteria -Pain intensity (VAS) between 4 and 8 while walking at the time of evaluation - The severity of OA on K-L classification system, 2 and 3 - Ability to walk more than 6 meters -Pain of one-sided of the knee (even if a person has bilateral arthritis, a knee that is painful will be included in the study) - knee pain for more than 3 months

##### Intervention groups

In the treatment group, the participants will receive the common physiotherapy of knee arthrosis with the Mulligan technique, and in the control group they will receive the common physiotherapy of the knee arthrosis with the Mulligan technique in an artificial form. After 10

sessions of physiotherapy, the intervention group will receive electrical muscle stimulation along with walking for 12 sessions.

##### Main outcome variables

Disability, spatiotemporal and kinematic characteristics of walking and pain

#### General information

##### Reason for update

Completion and termination of sampling and addition of outcome and the second phase of the intervention, which was approved by the ethics committee.

##### Acronym

##### IRCT registration information

IRCT registration number: **IRCT20190315043058N2**  
Registration date: **2024-01-09, 1402/10/19**  
Registration timing: **prospective**

Last update: **2025-12-31, 1404/10/10**

Update count: **1**

##### Registration date

2024-01-09, 1402/10/19

##### Registrant information

##### Name

Ismail Ebrahimi Takamjani

##### Name of organization / entity

##### Country

Iran (Islamic Republic of)

##### Phone

+98 21 2239 2426

##### Email address

ebrahimitakamjani.e@iums.ac.ir

##### Recruitment status

**Recruitment complete**

##### Funding source

**Expected recruitment start date**

2024-01-29, 1402/11/09

**Expected recruitment end date**

2024-08-30, 1403/06/09

**Actual recruitment start date**

2024-01-29, 1402/11/09

**Actual recruitment end date**

2024-10-30, 1403/08/09

**Trial completion date**

2024-12-29, 1403/10/09

**Scientific title**

The effects of Mobilization with movement technique and knee osteoarthritis conventional physiotherapy on disability, pain and spatiotemporal and kinematic characteristics of gait in individuals with knee osteoarthritis

**Public title**

The effects of Mobilization with movement technique and knee osteoarthritis conventional physiotherapy in knee osteoarthritis.

**Purpose**

Treatment

**Inclusion/Exclusion criteria****Inclusion criteria:**

1) Age 40 to 70 years 2) Diagnosis of knee tibiofemoral joint arthrosis based on the criteria of the American College of Rheumatology (knee pain for at least the last 3 months, joint crepitation and morning stiffness lasting 30 minutes or less) by a specialist physician. 3) Pain intensity based on visual pain scale between 4 and 8 while walking at the time of evaluation 4) The severity of arthritis should be 2 and 3 based on the Kellgren and Lawrence classification system. 5) Ability to walk more than 6 meters 6) One-sided arthritis pain of the tibiofemoral joint of the knee (even if a person has bilateral arthritis, a knee that is painful will be included in the study) 7) Having knee pain for more than 3 months

**Exclusion criteria:**

1) Any neurological problem that affects a person's walking, such as a stroke 2) Arthritis secondary to trauma or accident or fracture in the joint 3) Any surgery and fracture in any of the lower limbs in the last year 4) Injection of corticosteroid or hyaluronic acid or platelet-rich plasma in the last 6 months 5) Any pain in the upper or lower joints 6) Heart problems with medium and high risk, such as uncontrolled angina or heart infarction in the last month 7) Inflammation and infection in the knee 8) Any prohibition to manual treatments such as severe osteoporosis or unstable joint 9) Cognitive problem 10) Receiving physiotherapy intervention for knee arthritis in the last 12 months 11) Body mass index above 35 12) Using walking aids 13) Smoking 14) Systemic diseases such as rheumatoid arthritis

**Age**

From **40 years** old to **70 years** old

**Gender**

Both

**Phase**

N/A

**Groups that have been masked**

- Participant
- Outcome assessor
- Data analyser
- Data and Safety Monitoring Board

**Sample size**

Target sample size: **38**

Actual sample size reached: **38**

**Randomization (investigator's opinion)**

Randomized

**Randomization description**

Subjects will be divided into two control and treatment groups by the stratified randomization method, so the two groups will be balanced in terms of gender, and the process of this random division will be such that after creating blocks of 4 in each class (male gender and female) which consist of letters A and B, we will create a random sequence of these blocks using RANDOM NUMBER GENERATOR and the subjects will be placed in two control and treatment groups based on the order of reference by this sequence. The individual generating the randomization sequence will not participate in any other phase of the study.

**Blinding (investigator's opinion)**

Double blinded

**Blinding description**

In this study, the examiner or the person in charge of data collection and the one who evaluates the outcome and the subject or the participant will not know how the groups are divided. The examiner will not be present at the stage of dividing the subjects into treatment and control groups and the rest of the study stages. Also, the treatment that the subjects will receive in both groups will be apparently and look the same. In the treatment group, the subjects will receive preparatory treatments, therapeutic exercises and manual treatments in the form of the Mulligan technique, and in the control group, the subjects will receive preparatory treatments, therapeutic exercises and manual treatments in the form of the Mulligan technique in a simulated form; The method of performing this technique will be in such a way that no force will be applied to the place of handing after correct handing.

**Placebo**

Used

**Assignment**

Factorial

**Other design features****Secondary Ids**

empty

**Ethics committees****1****Ethics committee****Name of ethics committee**

Ethics committee of Iran University of Medical Sciences

**Street address**

Iran University of Medical Sciences, Hemet Highway,  
next to Milad Tower, Tehran

**City**

Tehran

**Province**

Tehran

**Postal code**

1341853347

**Approval date**

2023-10-23, 1402/08/01

**Ethics committee reference number**

IR.IUMS.REC.1402.665

**2****Ethics committee****Name of ethics committee**

Ethics committee of Iran University of Medical

**Street address**

Iran University of Medical Sciences, Hemet Highway,  
next to Milad Tower, Tehran

**City**

Tehran

**Province**

Tehran

**Postal code**

1341853347

**Approval date**

2023-10-29, 1402/08/07

**Ethics committee reference number**

IR.IUMS.REC.1402.668

**Health conditions studied****1****Description of health condition studied**

Knee Osteoarthritis

**ICD-10 code****ICD-10 code description****Primary outcomes****1****Description**

Disability score in Western Ontario and McMaster  
Universities Osteoarthritis Index questionnaire

**Timepoint**

Measurement of Western Ontario and McMaster  
Universities Osteoarthritis Index questionnaire score at  
the beginning of the study and after 10 sessions of  
physiotherapy treatment

**Method of measurement**

Western Ontario and McMaster Universities Osteoarthritis  
Index

**Secondary outcomes****1****Description**

Spatiotemporal and Kinematic characteristics of gait

**Timepoint**

Measurement of spatio-temporal and kinematic  
parameters of walking at the beginning of the study and  
after 10 sessions of physiotherapy treatment

**Method of measurement**

Vicon Motion Capture System

**2****Description**

Intensity of pain

**Timepoint**

Measurement of intensity of pain at the beginning of the  
study and after 10 sessions of physiotherapy treatment

**Method of measurement**

Visual Analogue Scale

**3****Description**

Characteristics of electromyographic activity of muscles  
during gait, sit to stand and static balance

**Timepoint**

Measurement of variables at the beginning of the study  
and after 10 sessions of physiotherapy treatment and  
after 12 sessions of home-based walking training.

**Method of measurement**

Kistler force plates and Myon electromyography system

**Intervention groups****1****Description**

Intervention group: Intervention group: The participants  
in this group will receive 10 treatment sessions during 4  
weeks with at least one day between sessions by a  
therapist who is proficient in the Mulligan technique and  
regular physical therapy. The treatment in each group  
lasts for 60 minutes and includes 3 phases of  
preparation, exercise therapy and manual treatments.  
The first phase, preparation: in this phase, subjects will  
receive high-frequency electrical nerve surface  
stimulation for 20 minutes and infra red for 15 minutes. 4  
electrodes are placed inside wet pads around the knee  
joint (two electrodes on the medial of the knee above  
and below the joint line and two electrodes on the lateral  
of the knee above and below the joint line) and will be  
tightened by the strap. The frequency and pulse duration  
used will be 110 Hz and 50 microseconds, respectively.  
The second phase, exercise therapy: The exercise  
therapy for these people will be in the form of  
strengthening exercises, stretching exercises, and range  
of motion exercises. Strengthening exercises include  
isotonic contraction with the help of varying leg weights  
of the knee extensor and hip abductor muscles,  
respectively, in sitting positions on the edge of the bed  
and lying on the side while the painful knee is elevated,  
with a frequency of 3 sessions per week according to the

ability of the subject with The intensity of 60% to 80% of 1RM will include 2 to 4 sets of 8 to 15 sets, and between each set we will allocate 30 to 60 seconds of rest time to the subjects. We will use stretching exercises and active range exercises for muscle shortening or reducing the range of motion that disrupts the person's performance. Stretching exercises will be performed for 30 seconds in 3 sets of 10 for the biceps and hamstring muscles, and active and passive range of motion exercises will be performed in the form of movement from fully bent knee to fully straightened knee in 3 sets of 10. will get The third phase, manual treatments: At first, we will perform these exercises while lying on the back with the knees slightly bent. If the subject does not have pain in bending or straightening the knee while lying on the back, we will perform this technique for the subject while standing. gave Also, if after a few treatment sessions, the examiner does not have pain in bending and straightening his knee while lying on his back, we will follow the technique while standing. The treatment process will be like this, lying on the back with a slightly bent knee, we will ask the subject to actively bend and straighten the knee, then after placing hands around the knee joint, the subject's leg will be internally rotated, respectively. We will perform external rotation, internal glide, external glide, anterior glide and posterior glide and at the same time we will ask the subject to move his knee in the same painful direction and if the active movement of the subject that was painful before with any of the rotations or glides It was done without pain, we will use the same rotation or glide to continue the treatment. This technique should be performed without pain or with minimal pain, and if the pain decreases but does not reach zero, the therapist will reach a completely pain-free state by increasing or decreasing the applied pressure or applying additional pressure at the end of the range. This technique will be performed in the first session as a set of 6 and in the following sessions, according to the subject's ability, as 3 to 5 sets of 6 to 10. To perform the technique in a standing position, the subject will place the sole of the foot whose knee is painful on a stool at a height of 30 cm ; Then we will take active knee bending and straightening movements by applying internal and external rotations, internal and external glides, and anterior and posterior glides to the subject's leg. If the rotations do not hurt, we will continue the painful movement with the same rotation or glide for the subject. Secondary intervention description: After completion 10 sessions of the physiotherapy intervention, participants will be re-randomized, such that an equal number of participants from each group will be allocated to the intervention and control groups. In this study, the knee joint with more severe symptoms will be selected for assessment of muscle activation and fitting of the unloader knee brace. In cases where symptom severity is equal in both knees, the dominant knee will be selected. Both groups will use the brace daily for 1 to 8 hours and will perform twelve 30-minute walking sessions over a four-week period. The control group will use the unloader knee brace without electrical stimulation, whereas the NMES group will receive neuromuscular electrical stimulation concurrently with

gait during walking. Stimulation intensity will be adjusted based on patient feedback up to the maximum tolerable level, such that painless muscle contractions are elicited, and participants will be encouraged to gradually increase the stimulation intensity. Electrode placement will be instructed by a physiotherapist, and patient adherence will be monitored through weekly follow-ups and adherence forms.

### **Category**

Treatment - Other

## **2**

### **Description**

Control group: Intervention group: Intervention group: The participants in this group will receive 10 treatment sessions during 4 weeks with at least one day between sessions by a therapist who is proficient in the Mulligan technique and regular physical therapy. The treatment in each group lasts for 60 minutes and includes 3 phases of preparation, exercise therapy and artificial manual treatments . The first phase, preparation: in this phase, subjects will receive high-frequency electrical nerve surface stimulation for 20 minutes and infra red for 15 minutes. 4 electrodes are placed inside wet pads around the knee joint (two electrodes on the medial of the knee above and below the joint line and two electrodes on the lateral of the knee above and below the joint line) and will be tightened by the strap. The frequency and pulse duration used will be 110 Hz and 50 microseconds, respectively. The second phase, exercise therapy: The exercise therapy for these people will be in the form of strengthening exercises, stretching exercises, and range of motion exercises. Strengthening exercises include isotonic contraction with the help of varying leg weights of the knee extensor and hip abductor muscles, respectively, in sitting positions on the edge of the bed and lying on the side while the painful knee is elevated, with a frequency of 3 sessions per week according to the ability of the subject with The intensity of 60% to 80% of 1RM will include 2 to 4 sets of 8 to 15 sets, and between each set we will allocate 30 to 60 seconds of rest time to the subjects. We will use stretching exercises and active range exercises for muscle shortening or reducing the range of motion that disrupts the person's performance. Stretching exercises will be performed for 30 seconds in 3 sets of 10 for the biceps and hamstring muscles, and active and passive range of motion exercises will be performed in the form of movement from fully bent knee to fully straightened knee in 3 sets of 10. The third phase, manual treatments: after the usual physical therapy exercises, the Mulligan technique will be performed in an artificial way for these patients in such a way that hands will be placed around the joint and the subject will be asked to bend and straighten his knee without applying glide or pressure, and we will perform this operation in 3 to 5 sets of 6 to 10, and we will give the patient a 60-second rest between each set. All the mentioned steps, including the evaluation to find the glide or rotation with pain-free movement, as well as the part of performing the technique for the treatment, for the implementation of this technique in the treatment group, will be done exactly for the control group, and as it was said, the only

difference will be this. It was that the glide or rotation in question will not be done in real way and no force will be applied and it will only be manual.

**Category**

Treatment - Other

**Recruitment centers****1****Recruitment center****Name of recruitment center**

Firoozgar general Hospital

**Full name of responsible person**

Hanieh Javan

**Street address**

Firoozgar Hospital, Beh Afarin St., Karim Khan Zand Blvd., Tehran, Iran

**City**

Tehran

**Province**

Tehran

**Postal code**

۱۵۹۳۷۴۸۷۱۱

**Phone**

+98 21 8214 1000

**Email**

firoozgarhospital1@gmail.com

**Sponsors / Funding sources****1****Sponsor****Name of organization / entity**

Iran University of Medical Sciences

**Full name of responsible person**

Dr.Reza falak

**Street address**

Office of Vice P Iran University of Medical Sciences (IUMS), Shahid Hemmat Highway, Tehran 14496-14535, IRAN

**City**

Tehran

**Province**

Tehran

**Postal code**

۱۴۴۹۶۱۴۵۳۵

**Phone**

+98 21 86701

**Email**

info@iums.ac.ir

**Grant name****Grant code / Reference number****Is the source of funding the same sponsor organization/entity?**

Yes

**Title of funding source**

Iran University of Medical Sciences

**Proportion provided by this source**

100

**Public or private sector**

Public

**Domestic or foreign origin**

Domestic

**Category of foreign source of funding**

empty

**Country of origin****Type of organization providing the funding**

Academic

**Person responsible for general inquiries****Contact****Name of organization / entity**

Iran University of Medical Sciences

**Full name of responsible person**

Hanieh Javan

**Position**

student

**Latest degree**

Bachelor

**Other areas of specialty/work**

Physiotherapy

**Street address**

Hanieh Javan.Unit 7, No 39, Alidoosti St, Ahaniamineh Ave, Tehran Pars

**City**

Tehran

**Province**

Tehran

**Postal code**

1653874986

**Phone**

+98 21 7788 9173

**Email**

javan.h@iums.ac.ir

**Person responsible for scientific inquiries****Contact****Name of organization / entity**

Iran University of Medical Sciences

**Full name of responsible person**

Dr. Esmail Ebrahimi Takamjani

**Position**

Associate professor

**Latest degree**

Ph.D.

**Other areas of specialty/work**

Physiotherapy

**Street address**

Faculty of Rehabilitation Sciences, Iran University of Medical Sciences, Shah Nazari St., Maddkaran St., Mader Square, Mirdamad, Tehran

**City**

Tehran

**Province**

Tehran

**Postal code**

۱۳۴۸۷ ۱۵۴۵۹

**Phone**

+98 21 2222 8051

**Email**

ebrahimitakamjani.e@iums.ac.ir

+98 21 2222 8051

**Email**

ebrahimitakamjani.e@iums.ac.ir

**Person responsible for updating data****Contact****Name of organization / entity**

Iran University of Medical Sciences

**Full name of responsible person**

Dr. Esmail Ebrahimi Takamjani

**Position**

Associate professor

**Latest degree**

Ph.D.

**Other areas of specialty/work**

Physiotherapy

**Street address**

Faculty of Rehabilitation Sciences, Iran University of  
Medical Sciences, Shah Nazari St., Maddkaran St.,  
Mader Square, Mirdamad, Tehran

**City**

Tehran

**Province**

Tehran

**Postal code**

۱۳۴۸۷ ۱۵۴۵۹

**Phone****Sharing plan****Deidentified Individual Participant Data Set (IPD)**

Undecided - It is not yet known if there will be a plan to make this available

**Study Protocol**

Undecided - It is not yet known if there will be a plan to make this available

**Statistical Analysis Plan**

Undecided - It is not yet known if there will be a plan to make this available

**Informed Consent Form**

Undecided - It is not yet known if there will be a plan to make this available

**Clinical Study Report**

Undecided - It is not yet known if there will be a plan to make this available

**Analytic Code**

Undecided - It is not yet known if there will be a plan to make this available

**Data Dictionary**

Undecided - It is not yet known if there will be a plan to make this available