

Clinical Trial Protocol

Iranian Registry of Clinical Trials

10 Jun 2026

Comparison of the effectiveness of vaginal dexamethasone and placebo in preparing the cervix for induction of labor in primiparous pregnant women

Protocol summary

Study aim

Determining the effectiveness of vaginal dexamethasone and placebo in preparing the cervix for induction of labor in primiparous pregnant women referred to Ali-Ebne-Abitaleb hospital in Zahedan during 2025-2026

Design

Clinical trial with intervention and control groups, triple-blind, randomized block design (4-way permutation-stratified), phase 2-3, sample size 100

Settings and conduct

Triple-blind randomized clinical trial in Ali ibn Abi Talib Hospital, Zahedan Number of participants: 100 patients (two groups of 50 each) Groups: Intervention: Vaginal dexamethasone tablet (Tablet A) Comparison: Vaginal placebo (Tablet B) Blinding: The patients, researchers, and research staff are all blinded.

Participants/Inclusion and exclusion criteria

All pregnant mothers referred to the Women's Triage Unit of Ali Ibn Abi Taleb Hospital, Zahedan. Inclusion criteria: Primiparous pregnant mother, singleton pregnancy, age between 18 and 35 years Exclusion criteria: The pregnant woman has used any hormones or medications during her pregnancy. If the pregnant woman has previously referred with immobility/decreased fetal movements or a history of maternal bleeding

Intervention groups

The intervention group consisted of nulliparous pregnant women who received vaginal dexamethasone tablets (Tablet A). The comparison group consisted of nulliparous pregnant women who received vaginal placebo (Tablet B).

Main outcome variables

Bishop score; Apgar score at first minute of the newborn; Apgar score at fifth minute of the newborn

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20250118064418N1**

Registration date: **2025-03-01, 1403/12/11**

Registration timing: **registered_while_recruiting**

Last update: **2025-03-01, 1403/12/11**

Update count: **0**

Registration date

2025-03-01, 1403/12/11

Registrant information

Name

setareh jami

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 915 763 6126

Email address

setarehjami9111@gmail.com

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2025-02-08, 1403/11/20

Expected recruitment end date

2025-06-10, 1404/03/20

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Comparison of the effectiveness of vaginal dexamethasone and placebo in preparing the cervix for induction of labor in primiparous pregnant women

Public title

Efficacy of vaginal Dexamethasone in cervical preparation

Purpose

Treatment

Inclusion/Exclusion criteria

Inclusion criteria:

Primiparous pregnant mother singleton pregnancy full pregnancy (37 weeks and 6 days and more) age between 18 and 35 years normal body mass index 18.5 to 24.9 the fetus must be in cephalic position Bishop's score ≤ 4 estimated fetal weight of 2500-4000 grams the gynecologist and obstetrician make the patient a candidate for termination of pregnancy

Exclusion criteria:

If the pregnant woman use any hormones or herbal medicines (traditional medicines), herbal/natural products, or conventional/prescribed medicines during her pregnancy Those with the following pre-existing conditions, including diabetes, high blood pressure, a history of obstetric complications such as preeclampsia or eclampsia, or HELLP syndrome (hemolysis, elevated liver enzymes, and low platelets) Those with a previous presentation with fetal inactivity/decreased fetal movements Those with a previous presentation with a history of maternal hemorrhage

Age

From **18 years** old to **35 years** old

Gender

Female

Phase

2-3

Groups that have been masked

- Participant
- Care provider
- Investigator
- Outcome assessor
- Data analyser

Sample size

Target sample size: **100**

Randomization (investigator's opinion)

Randomized

Randomization description

1. Randomization method and description of each method Type of randomization: Block randomization with blocks of four was used in this study 2. Randomization unit Individual randomization: Subjects are entered into the study individually based on patient referrals and are assigned to intervention and control groups through block randomization. 3. Stratified randomization layers If it is necessary to control the effect of intervening variables (such as age, sex, and underlying diseases), stratified randomization is used to perform randomization in each layer in such a way that the balance between groups is maintained regarding important variables. 4. Randomization tool and method

of constructing random sequences: It will be using SPSS software. Number of samples: Finally, 50 subjects were placed in the intervention group and 50 subjects in the control group to maintain balance in the sample size.

Blinding (investigator's opinion)

Triple blinded

Blinding description

After enrollment, patients, investigators, and research staff are blinded. Dexamethasone and placebo tablets of the same shape and size, labeled Tablet A and Tablet B, are dispensed by a non-interventionist in the study. Research staff are blinded to the allocation of study drug tablets.

Placebo

Used

Assignment

Parallel

Other design features

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics committee of Zahedan University of Medical Sciences

Street address

Khalij -e- Fars Blvd

City

Zahedan

Province

Sistan-va-Balouchestan

Postal code

9816743175

Approval date

2024-10-16, 1403/07/25

Ethics committee reference number

IR.ZAUMS.REC.1403.287

Health conditions studied

1

Description of health condition studied

Induction of labor with dexamethasone

ICD-10 code

ICD-10 code description

Primary outcomes

1

Description

Cervical preparation rate for labor induction in primiparous pregnant mothers

Timepoint

During labor, the first minute, and the fifth minute after

delivery

Method of measurement

Cervical preparation dosage form for labor induction

Secondary outcomes

1

Description

Newborn Apgar score

Timepoint

The first and fifth minutes after birth

Method of measurement

The Apgar score, also known as APGAR, is an acronym for 5 criteria that are each examined separately: Skin color (Appearance), Pulse or heart rate (Pulse), Reflex reactions (Grimace), Muscle tone or consistency (Activity), and Respiration.

2

Description

Bishop Score

Timepoint

The initial visit of the pregnant woman and entry into the study and during delivery

Method of measurement

Based on 5 components of dilation, effacement, consistency, cervical position, and organ position, demonstrated through pelvic examination, each of which is given a score from 0 to 3.

Intervention groups

1

Description

Intervention group: Dexamethasone tablets labeled as Tablet A are delivered by a non-interventionist in the study. Dexamethasone is administered in the amount of 8 mg in the form of 2 Dexamethasone tablets (4 mg, from a single company (Obeidi Company) and specified) every 6 hours and vaginally in group A. According to the study protocol, an obstetrician and gynecologist will prescribe the tablets. Vaginal tablets are given to primiparous patients from the onset of labor pains and it is recommended to use 2 tablets every 6 hours and vaginally. If possible, do not move for at least 2 hours after using the tablet. Also, to ensure the use of the tablet, a daily registration form is given to the participants to record each time the tablet is used.

Category

Treatment - Other

2

Description

Control group: Placebo labeled as pill B is delivered by a non-interventionist in the study. It is used as 2 pills (completely identical to the intervention drug) every 6 hours and vaginally. According to the study protocol, an obstetrician-gynecologist will prescribe the pill. Vaginal

pills are given to primiparous patients from the onset of labor pains and it is recommended to use 2 pills every 6 hours and vaginally. If possible, do not move for at least 2 hours after using the pill. Also, to ensure the use of the pill, participants are given a daily registration form to record each time the pill is used.

Category

Treatment - Other

Recruitment centers

1

Recruitment center

Name of recruitment center

Women's triage at Ali Ibn Abi Taleb Hospital in Zahedan.

Full name of responsible person

Setareh Jami

Street address

Khalij -e- Fars Blvd

City

Zahedan

Province

Sistan-va-Balouchestan

Postal code

9816743175

Phone

+98 915 763 6126

Email

seterahjami9111@gmail.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Zahedan University of Medical Sciences

Full name of responsible person

Ebrahim Kord

Street address

Khalij -e- Fars Blvd

City

Zahedan

Province

Sistan-va-Balouchestan

Postal code

9816743175

Phone

+98 54 3337 2117

Email

e.kord58@gmail.com

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Zahedan University of Medical Sciences

Proportion provided by this source

100

Public or private sector

Public

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin**Type of organization providing the funding**

Academic

Person responsible for general inquiries**Contact****Name of organization / entity**

Zahedan University of Medical Sciences

Full name of responsible person

Setareh Jami

Position

Resident

Latest degree

Medical doctor

Other areas of specialty/work

Gynecology and Obstetrics

Street address

Khalij -e- Fars Blvd

City

Zahedan

Province

Sistan-va-Balouchestan

Postal code

9816743175

Phone

+98 915 763 6126

Email

setarehjami9111@gmail.com

Person responsible for scientific inquiries**Contact****Name of organization / entity**

Zahedan University of Medical Sciences

Full name of responsible person

Setareh Jami

Position

Resident

Latest degree

Medical doctor

Other areas of specialty/work

Gynecology and Obstetrics

Street address

Khalij -e- Fars Blvd

City

Zahedan

Province

Sistan-va-Balouchestan

Postal code

9816743175

Phone

+98 915 763 6126

Email

setarehjami9111@gmail.com

Person responsible for updating data**Contact****Name of organization / entity**

Zahedan University of Medical Sciences

Full name of responsible person

Setareh jami

Position

resident

Latest degree

Medical doctor

Other areas of specialty/work

Gynecology and Obstetrics

Street address

Khalij -e- Fars Blvd

City

zahedan

Province

Sistan-va-Balouchestan

Postal code

9816743175

Phone

+98 915 763 6126

Email

setarehjami9111@gmail.com

Sharing plan**Deidentified Individual Participant Data Set (IPD)**

Yes - There is a plan to make this available

Study Protocol

Yes - There is a plan to make this available

Statistical Analysis Plan

Yes - There is a plan to make this available

Informed Consent Form

Yes - There is a plan to make this available

Clinical Study Report

Yes - There is a plan to make this available

Analytic Code

Yes - There is a plan to make this available

Data Dictionary

Yes - There is a plan to make this available

Title and more details about the data/document

Information about the main outcome

When the data will become available and for how long

Access period starts 6 months after results are published.

To whom data/document is available

Researchers working in academic and scientific institutions

Under which criteria data/document could be used

The use of data for research purposes will be permitted. A written request from the requesting person and a guarantee of confidentiality of the information are required.

From where data/document is obtainable

Email: setareh jami setarehjami9111@gmail.com

What processes are involved for a request to access data/document

Submitting a request - Reviewing the request and complying with scientific and ethical principles -

