

Clinical Trial Protocol

Iranian Registry of Clinical Trials

29 Aug 2026

Comparison of the effect of pioglitazone and Gloranta on metabolic factors of patients with non-alcoholic fatty liver disease and type 2 diabetes

Protocol summary

Study aim

Comparison of the effects of pioglitazone and Glurenta on metabolic factors in patients with non-alcoholic fatty liver disease and type 2 diabetes referred to Clinic

Design

Single-blind, non-controlled phase 4 clinical trial on 72 patients

Settings and conduct

In this single-blind trial, the researchers involved in assessing the outcomes are unaware of the type of intervention that the patients receive. The study uses an independent person to collect the data, who has no connection to the intervention, and after the data is collected, the results are analyzed. Patients are assigned to groups using random blocks after obtaining informed consent. Also, in this study, no additional testing or costs are imposed on patients in addition to the diagnostic and treatment process. 72 patients with diabetes and non-alcoholic fatty liver disease were divided into two groups of 36, one group receiving pioglitazone 15 mg daily and the other group receiving 10.5 mg daily. The dose of 10.5 mg of gloranta is an effective dose. After grouping, pioglitazone and gloranta are administered orally once a day for 6 months.

Participants/Inclusion and exclusion criteria

Patients aged 18 to 75 years with non-alcoholic fatty liver disease and type 2 diabetes No history of alcohol consumption (20 grams per day for women and 30 grams per day for men)

Intervention groups

72 patients with diabetes and non-alcoholic fatty liver disease were divided into two groups of 36, one group receiving pioglitazone 15 mg daily and the other group receiving 10.5 mg daily. The dose of 10.5 mg daily is an effective dose. After grouping, pioglitazone and 10.5 mg daily are administered orally for 6 months.

Main outcome variables

Lipid profile; BMI; Weight; Blood pressure; Liver enzymes; Fatty liver level; Blood sugar; Platelets; ESR, CRP

General information

Reason for update

Acronym

COEOPAG

IRCT registration information

IRCT registration number: **IRCT20250919067293N1**

Registration date: **2025-09-29, 1404/07/07**

Registration timing: **prospective**

Last update: **2025-09-29, 1404/07/07**

Update count: **0**

Registration date

2025-09-29, 1404/07/07

Registrant information

Name

Fatemeh Bahrapour

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 21 5517 3894

Email address

nemesisgrace@googlemail.com

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2025-10-17, 1404/07/25

Expected recruitment end date

2026-04-20, 1405/01/31

Actual recruitment start date

empty

Actual recruitment end date
empty

Trial completion date
empty

Scientific title
Comparison of the effect of pioglitazone and Gloranta on metabolic factors of patients with non-alcoholic fatty liver disease and type 2 diabetes

Public title
Comparison of the effect of pioglitazone and Gloranta on metabolic factors of patients with non-alcoholic fatty liver disease and type 2 diabetes

Purpose
Treatment

Inclusion/Exclusion criteria
Inclusion criteria:
Patients aged 18 to 75 years with non-alcoholic fatty liver disease and type 2 diabetes No history of alcohol consumption (20 grams per day for women and 30 grams per day for men) No history of cardiovascular events in the past 3 months Not pregnant or breastfeeding; active cancer or history of cancer treatment in the past 2 years; body mass index above 40 kg/m2.
Exclusion criteria:
Active or chronic hepatitis; cirrhosis and biliary disease; heart failure, renal dysfunction (eGFR <45 ml/min/1.73 m2) during the study Taking medications associated with fatty liver nonsteroidal anti-inflammatory drugs (NSAIDs) (amiodarone, tamoxifen, sodium valproate, corticosteroids, methotrexate). Simultaneous use of both drugs, Glurenta and Empagliflozin

Age
From **18 years** old to **75 years** old

Gender
Both

Phase
4

Groups that have been masked

- Data analyser

Sample size
Target sample size: **72**
More than 1 sample in each individual
Number of samples in each individual: **2**
Blood sample and ultrasound

Randomization (investigator's opinion)
Not randomized

Randomization description

Blinding (investigator's opinion)
Single blinded

Blinding description
In this single-blind trial, the researchers involved in assessing the outcomes are unaware of the type of intervention that the patients receive. The study uses an independent person to collect the data who has no connection to the intervention, and after the data is collected, the results will be analyzed.

Placebo

Not used

Assignment
Parallel

Other design features
no

Secondary Ids

empty

Ethics committees

1

Ethics committee

Name of ethics committee

Ethics Committee of Baqiyatullah University of Medical Sciences

Street address

No. 24, Hakimian Alley, Badakhshan Kamali Street, Abuzar Street

City

Tehran

Province

Tehran

Postal code

1363935191

Approval date

2025-08-06, 1404/05/15

Ethics committee reference number

IR.BMSU.REC.1404.015

Health conditions studied

1

Description of health condition studied

Diabetes mellitus

ICD-10 code

E08

ICD-10 code description

Diabetes mellitus due to underlying condition

2

Description of health condition studied

Non-alcoholic fatty liver

ICD-10 code

K76.0

ICD-10 code description

Fatty (change of) liver, not elsewhere classified

Primary outcomes

1

Description

Liver enzymes

Timepoint

Measurement of liver enzymes at the beginning of the study before the start of the intervention and 6 months after the start of Glurenta or pioglitazone

Method of measurement

A blood sample

2**Description**

blood pressure

Timepoint

At the beginning of the study before the start of the intervention and 6 months after starting Gloreanta or pioglitazone

Method of measurement

Blood pressure measuring device

3**Description**

Body mass index of individuals

Timepoint

At the beginning of the study before the start of the intervention and 6 months after starting Gloreanta or pioglitazone

Method of measurement

Scales and meters

4**Description**

, LDL, HDL, Total cholesterol, Triglycerides

Timepoint

At the beginning of the study before the start of the intervention and 6 months after starting Gloreanta or pioglitazone

Method of measurement

blood sample

5**Description**

Fatty liver

Timepoint

At the beginning of the study before the start of the intervention and 6 months after starting Gloreanta or pioglitazone

Method of measurement

The amount of liver fat is determined by ultrasound and fibroscan and based on the interpretation of the radiologist.

6**Description**

(FBS, BS2hpp, Hba1c)

Timepoint

At the beginning of the study before the start of the intervention and 6 months after starting Gloreanta or pioglitazone

Method of measurement

blood sample

7**Description**

platelet

Timepoint

At the beginning of the study before the start of the intervention and 6 months after starting Gloreanta or pioglitazone

Method of measurement

blood sample

Secondary outcomes

empty

Intervention groups**1****Description**

Intervention group: Intervention group: 72 people with diabetes and non-alcoholic fatty liver disease were divided into two groups of 36 people: one group received pioglitazone 15 mg daily and the other group received 10.5 mg daily. The dose of Glurenta is 10.5 mg. An effective dose. Glurenta is administered orally once a day for 6 months. Then a blood sample is taken. Fasting blood glucose levels after 8 hours of fasting, blood levels of alanine aminotransferase (ALT), triglycerides (TG), free fatty acids (FFA) and CBC, ultrasound and cases requiring fibroscan are measured.

Category

Treatment - Drugs

2**Description**

Intervention group: Intervention group: 72 patients with diabetes and non-alcoholic fatty liver disease were divided into two groups of 36 people: one group received pioglitazone 15 mg daily and the other group received 10.5 mg daily. After grouping, pioglitazone was administered orally once a day for 6 months. Then a blood sample is taken. Fasting blood glucose levels after 8 hours of fasting, blood levels of alanine aminotransferase (ALT), triglycerides (TG), free fatty acids (FFA), and CBC, and ultrasound and in cases requiring fibroscan are measured.

Category

Treatment - Drugs

Recruitment centers**1****Recruitment center****Name of recruitment center**

Baqiyatullah Hospital Clinic

Full name of responsible person

Fatemeh Bahrampour

Street address

No. 24, Hakimian Alley, Badakhshan Kamali St., Abuzar St.

City

Tehran

Province
Tehran
Postal code
1363935191
Phone
+98 21 5517 3894
Email
Nemesisgrace@googlemail.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity
Bagheiat-allah University of Medical Sciences
Full name of responsible person
Mohammad javad Behzad nia
Street address
Mulla Sadra Street, Baqiyatallah Hospital
City
Tehran
Province
Tehran
Postal code
1363935191
Phone
+98 21 8126 7046
Email
bmcqomclinic@gmail.com
Grant name
Grant code / Reference number
Is the source of funding the same sponsor organization/entity?
Yes
Title of funding source
Bagheiat-allah University of Medical Sciences
Proportion provided by this source
1
Public or private sector
Public
Domestic or foreign origin
Domestic
Category of foreign source of funding
empty
Country of origin
Type of organization providing the funding
Academic

Person responsible for general inquiries

Contact

Name of organization / entity
Bagheiat-allah University of Medical Sciences
Full name of responsible person
Fatemeh Bahrapour
Position
resident
Latest degree
Medical doctor
Other areas of specialty/work
Internal Medicine

Street address
Abuzar Street, Bad Akhshan Kamali Street, No. 24,
Hakimian Alley
City
Tehran
Province
Tehran
Postal code
1363935191
Phone
+98 21 5517 3894
Email
Nemesisgrace@googlemail.com

Person responsible for scientific inquiries

Contact

Name of organization / entity
Bagheiat-allah University of Medical Sciences
Full name of responsible person
Fatemeh Bahrapour
Position
Resident
Latest degree
Medical doctor
Other areas of specialty/work
Internal Medicine
Street address
Abuzar Street, Bad Akhshan Kamali Street, No. 24,
Hakimian Alley
City
Tehran
Province
Tehran
Postal code
1363935191
Phone
+98 21 5517 3894
Email
Nemesisgrace@googlemail.com

Person responsible for updating data

Contact

Name of organization / entity
Bagheiat-allah University of Medical Sciences
Full name of responsible person
Fatemeh Bahrapour
Position
Resident
Latest degree
Medical doctor
Other areas of specialty/work
Internal Medicine
Street address
Abuzar Street, Bad Akhshan Kamali Street, No. 24,
Hakimian Alley
City
Tehran
Province
Tehran

Postal code

1363935191

Phone

+98 21 5517 3894

Email

Nemesisgrace@googlemail.com

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Undecided - It is not yet known if there will be a plan to make this available

Study Protocol

Undecided - It is not yet known if there will be a plan to make this available

Statistical Analysis Plan

Undecided - It is not yet known if there will be a plan to make this available

Informed Consent Form

Yes - There is a plan to make this available

Clinical Study Report

Undecided - It is not yet known if there will be a plan to

make this available

Analytic Code

Undecided - It is not yet known if there will be a plan to make this available

Data Dictionary

Undecided - It is not yet known if there will be a plan to make this available

Title and more details about the data/document

no

When the data will become available and for how long

6 months after results are published

To whom data/document is available

No body

Under which criteria data/document could be used

no

From where data/document is obtainable

no

What processes are involved for a request to access data/document

NO

Comments