

Clinical Trial Protocol

Iranian Registry of Clinical Trials

11 Sep 2026

Comparing the effect of home-based synchronous tele-exercise on balance, fall risk, postural abnormalities, quality of life, physical activity level, and functional status in hemodialysis patients

Protocol summary

Study aim

To investigate the effect of 12 weeks of home-based corrective exercises with and without remote synchronous supervision on balance, fall risk, postural abnormalities, quality of life, physical activity level, and functional status in hemodialysis patients

Design

Randomized controlled clinical trial; parallel-group design; single-blind (outcome assessor); simple randomization using sealed opaque envelopes; sample size of 60 participants (30 per group); study duration of 5 months.

Settings and conduct

The study will be conducted at the dialysis center of Khorshid Hospital, Isfahan. Patients will be randomly assigned to two groups after informed consent. Assessments of balance, fall risk, walking ability, postural abnormalities, quality of life, and physical activity will be performed before and after the intervention. The intervention group will perform home exercises on non-dialysis days with online supervision by a trainer, and the outcome assessor will be unaware of the patient grouping.

Participants/Inclusion and exclusion criteria

Inclusion criteria: Patients referred to Khorshid Hospital Dialysis Center; age 40-60 years; at least 3 months dialysis history; 3 times weekly hemodialysis; no recent heart attack/stroke; able to communicate; physician permission; exclusion: unstable cardiac status; active infection; unstable blood sugar; unable to exercise; severe musculoskeletal pain; shortness of breath at rest/daily activities

Intervention groups

Intervention group: 12 weeks, 3 sessions/week home exercises (warm-up 5 min, corrective/strength 15-30 min, cool-down 10 min) with synchronous online supervision via WhatsApp; Control group: same exercises

via printed booklet without supervision; both groups continue usual care

Main outcome variables

Balance (assessed by Berg Balance Scale); fall risk (assessed by Tinetti Performance-Oriented Mobility Assessment); forward head posture (assessed by craniovertebral angle from digital photography); kyphosis (assessed by spinal angle measurement using a flexible ruler); quality of life; physical activity level; functional status (assessed by six-minute walking test)

General information

Reason for update

Acronym

IRCT registration information

IRCT registration number: **IRCT20200819048461N5**

Registration date: **2026-08-08, 1405/05/17**

Registration timing: **prospective**

Last update: **2026-08-08, 1405/05/17**

Update count: **0**

Registration date

2026-08-08, 1405/05/17

Registrant information

Name

Mohammad ali Tabibi

Name of organization / entity

Country

Iran (Islamic Republic of)

Phone

+98 31 3230 1795

Email address

m.tabibi@ut.ac.ir

Recruitment status

Not yet recruiting

Funding source

Expected recruitment start date

2026-09-23, 1405/07/01

Expected recruitment end date

2027-02-20, 1405/12/01

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Comparing the effect of home-based synchronous tele-exercise on balance, fall risk, postural abnormalities, quality of life, physical activity level, and functional status in hemodialysis patients

Public title

The effect of home-based synchronous tele-exercise on balance, risk of fall, abnormality of posture, quality of life, level of physical activity and physical function in dialysis patients

Purpose

Treatment

Inclusion/Exclusion criteria**Inclusion criteria:**

Age between 40 and 60 years At least three months of dialysis history Receiving hemodialysis three times per week No heart attack and stroke in the last three months Ability to communicate with others Doctor's permission for exercise

Exclusion criteria:

Unstable cardiac status (angina, congestive heart failure, severe arterial stenosis, uncontrolled arrhythmias, etc.) Active infection or acute medical disease Individuals with medically diagnosed unstable blood glucose levels (fluctuations between hypoglycemia and hyperglycemia) Unable to exercise (lower limb amputation without prosthesis) Severe musculoskeletal pain at rest or minimal activity Shortness of breath at rest or with daily activities

Age

From **40 years** old to **60 years** old

Gender

Both

Phase

N/A

Groups that have been masked

- Outcome assessor

Sample size

Target sample size: **60**

Randomization (investigator's opinion)

Randomized

Randomization description

A simple random allocation sequence will be generated by an independent person using a random number generator software. Each allocation will be placed in a sequentially numbered, opaque, sealed envelope. After obtaining informed consent, the next envelope will be opened and the patient will be assigned to either the

intervention or control group according to its content. Therefore, neither the researcher nor the patient is aware of the group assignment until the envelope is opened.

Blinding (investigator's opinion)

Single blinded

Blinding description

The study is single-blind, and the collaborator who takes the functional tests from the patients and measures postural abnormalities with a flexible ruler and digital photography and completes the questionnaires will not be aware of the patient grouping

Placebo

Not used

Assignment

Parallel

Other design features

This study is a randomized controlled clinical trial with a parallel-group design and blinded outcome assessment. The unique feature of this trial is the comparison of home-based corrective exercise with synchronous online coach supervision in the intervention group versus the same exercise program without online supervision in the control group. Participants will be allocated using simple randomization with concealed allocation through opaque sealed envelopes.

Secondary Ids

empty

Ethics committees**1****Ethics committee****Name of ethics committee**

Ethics Committee of Isfahan University of Medical Sciences

Street address

Isfahan University of Medical Sciences, Hezar Jerib Street, Isfahan, Iran

City

Isfahan

Province

Isfahan

Postal code

8174673461

Approval date

2022-10-22, 1401/07/30

Ethics committee reference number

IR.ARI.MUI.REC.1401.209

Health conditions studied**1****Description of health condition studied**

End-stage renal disease requiring hemodialysis

ICD-10 code

N18.6

ICD-10 code description

End stage renal disease

Primary outcomes

1

Description

Balance

Timepoint

Before intervention; after 12 weeks

Method of measurement

Berg Balance Scale (BBS), a 14-item performance-based instrument (e.g., sit-to-stand, unsupported standing, 360° turn, single-leg stance), each item scored 0 (unable) to 4 (independent); total score range 0–56, with lower scores indicating greater balance impairment and fall risk. Administered by a trained, blinded assessor in a clinical setting over approximately 15–20 minutes.

2

Description

Fall risk

Timepoint

Before intervention; after 12 weeks

Method of measurement

Tinetti Performance-Oriented Mobility Assessment (POMA), comprising a balance subscale (9 items, max 16 points) and a gait subscale (7 items, max 12 points); total score 0–28, with <19 indicating high fall risk, 19–23 moderate risk, and ≥24 low risk. Administered via direct observation by an assessor blinded to group allocation.

3

Description

Forward head posture

Timepoint

Before intervention; after 12 weeks

Method of measurement

Craniovertebral angle (CVA) measured from a lateral-view digital photograph taken in natural standing posture; anatomical landmarks (spinous process of C7 and the tragus of the ear) are marked on the skin, and the angle between the horizontal line through the tragus and the line connecting the tragus to the C7 spinous process is calculated using AutoCAD (or equivalent angle-measurement software); a smaller angle indicates greater forward head posture severity (normal typically >50°).

4

Description

Kyphosis

Timepoint

Before intervention; after 12 weeks

Method of measurement

Thoracic kyphosis angle measured using a Flexible Ruler; the ruler is molded over the T1–T12 spinous processes with the patient in natural standing posture and the spinal curve traced onto paper; the kyphosis angle is then calculated from the traced curve using the

trigonometric formula $\text{Kyphosis Angle} = \arctan(4H/L)$, where H is the height of the curve and L is the baseline length; higher values indicate greater kyphotic curvature.

5

Description

Quality of life

Timepoint

Before intervention; after 12 weeks

Method of measurement

Kidney Disease Quality of Life – Short Form (KDQOL-36), comprising the generic SF-12 core plus kidney disease-specific components (symptoms/problems, effects of kidney disease on daily life, burden of kidney disease); each domain is scored 0–100, with higher scores indicating better quality of life. Self-administered by the patient after the dialysis session, with assistance from the research collaborator if needed.

6

Description

Physical activity level

Timepoint

Before intervention; after 12 weeks

Method of measurement

Community Healthy Activities Model Program for Seniors (CHAMPS) questionnaire, covering approximately 41 physical and social activities; participants report frequency and duration of each activity over the past week; output is expressed as estimated weekly caloric expenditure (kcal/kg/week) and weekly hours of moderate-to-vigorous activity, with higher scores indicating greater physical activity. Administered as an interviewer-based questionnaire by a blinded research collaborator.

7

Description

Functional status

Timepoint

Before intervention; after 12 weeks

Method of measurement

Six-Minute Walk Test (6MWT), conducted per the American Thoracic Society (ATS) standard protocol; the patient walks at maximal comfortable pace along a flat 30-meter corridor for 6 minutes, and the total distance covered (in meters) is recorded; a greater distance indicates better functional exercise capacity. Performed on a non-dialysis day by a blinded assessor.

Secondary outcomes

1

Description

Hospitalization

Timepoint

During the study (5 months)

Method of measurement

Number and cause of hospital admissions during the 5-month study period, recorded via a researcher-made checklist based on medical record review and patient self-report at each dialysis session.

2

Description

Mortality

Timepoint

During the study (5 months)

Method of measurement

Recording of mortality cases and cause of death during the 5-month study period, via a researcher-made checklist based on review of the dialysis center's medical records.

Intervention groups

1

Description

Intervention group: Experimental group (with coach supervision): 12 weeks; 3 sessions/week exercise program at home on non-dialysis days with synchronous online supervision by coach via WhatsApp. The exercise program includes: general warm-up (5 minutes), corrective and strength exercises (15 minutes) and general cool-down (10 minutes), with the main exercise body starting at 15 minutes and gradually increasing by adding sets and repetitions to 30 minutes over three months. After each exercise, a 30 to 60 second rest is given to the patient.

Category

Rehabilitation

2

Description

Control group (without supervision): Participants receive the same home exercise program in a printed booklet but perform the exercises independently without synchronous online supervision. Both groups continue to receive usual medical care.

Category

Placebo

Recruitment centers

1

Recruitment center

Name of recruitment center

Khorshid Hospital Dialysis Center

Full name of responsible person

Shahrzad Shahidi

Street address

Dialysis Ward, Khorshid Hospital, Governorate Street

City

Isfahan

Province

Isfahan

Postal code

8165837311

Phone

+98 31 3222 2127

Email

shahidi_shahrzad@yahoo.com

Sponsors / Funding sources

1

Sponsor

Name of organization / entity

Pardis Specialized Wellness Institute

Full name of responsible person

Dr Mohamad Ali Tabibi

Street address

71 Alley. Jey Street

City

Isfahan

Province

Isfahan

Postal code

8156135414

Phone

+98 913 318 4624

Email

m.tabibi@ut.ac.ir

Grant name

Grant code / Reference number

Is the source of funding the same sponsor organization/entity?

Yes

Title of funding source

Pardis Specialized Wellness Institute

Proportion provided by this source

100

Public or private sector

Private

Domestic or foreign origin

Domestic

Category of foreign source of funding

empty

Country of origin

Type of organization providing the funding

Persons

Person responsible for general inquiries

Contact

Name of organization / entity

Pardis Specialized Wellness Institute

Full name of responsible person

Nasrin Salimian

Position

Head of Research and Statistics Department, Pardis Wellness Clinic Institute

Latest degree

Master

Other areas of specialty/work

Biostatistics

Street address

71 Alley. Jey Street
City
Isfahan
Province
Isfahan
Postal code
8156135414
Phone
0098 31 335252020
Email
statisrin@gmail.com

Person responsible for scientific inquiries

Contact

Name of organization / entity
Pardis Specialized Wellness Institute
Full name of responsible person
Dr. Mohammad Ali Tabibi
Position
Head of Pardis Specialized Wellness Institute
Latest degree
Ph.D.
Other areas of specialty/work
Physiology
Street address
71 alley. Jey street
City
Isfahan
Province
Isfahan
Postal code
8174757536
Phone
0098 31 335252020
Email
m.tabibi@ut.ac.ir

Person responsible for updating data

Contact

Name of organization / entity
Pardis Specialized Wellness Institute
Full name of responsible person
Dr. Mohammad Ali Tabibi
Position
Head of Pardis Specialized Wellness Institute
Latest degree
Ph.D.

Other areas of specialty/work

Physiology
Street address
71 alley. Jey street
City
Isfahan
Province
Isfahan
Postal code
8174757536
Phone
+98 31335252020
Email
m.tabibi@ut.ac.ir
Web page address

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Yes - There is a plan to make this available

Study Protocol

Yes - There is a plan to make this available

Statistical Analysis Plan

Yes - There is a plan to make this available

Informed Consent Form

Yes - There is a plan to make this available

Clinical Study Report

Yes - There is a plan to make this available

Analytic Code

Yes - There is a plan to make this available

Data Dictionary

Yes - There is a plan to make this available

Title and more details about the data/document

The results of the study group are made available

When the data will become available and for how long

After publication of the main results

To whom data/document is available

Only academics are allowed

Under which criteria data/document could be used

If the data is used for the purpose of research in the field of variables and providing new solutions, it can be used by mentioning the source

From where data/document is obtainable

Dr. Mohammad Ali Tabibi via email address
m.tabibi@ut.ac.ir

What processes are involved for a request to access data/document

It will be answered within one month after receiving the email

Comments