

Clinical Trial Protocol

Iranian Registry of Clinical Trials

27 Jul 2026

Comparison of Intraoperative Awareness with Dexmedetomidine-Propofol Infusion versus Intravenous Xylocaine-Propofol Infusion in Patients Undergoing Coronary Artery Bypass Grafting: A Randomized Controlled Trial

Protocol summary

Study aim

To compare the effectiveness of dexmedetomidine plus propofol infusion versus intravenous xylocaine (lidocaine) plus propofol infusion in preventing intraoperative awareness among patients undergoing elective coronary artery bypass grafting (CABG) under general anesthesia.

Design

RCT

Settings and conduct

National hospital lahore

Participants/Inclusion and exclusion criteria

Patients aged 30–75 years, of either gender. Scheduled for elective isolated CABG under general anesthesia. ASA physical status III–IV. LVEF $\geq 35\%$. Able to provide written informed consent and complete the Modified Brice Interview postoperatively. Exclusion Criteria Emergency or combined cardiac surgery, or previous cardiac surgery. Allergy or contraindication to dexmedetomidine, propofol, or lidocaine. Neurological or psychiatric disorders affecting awareness assessment. Chronic use of sedatives, opioids, alcohol, or recreational drugs. Severe hepatic or renal disease, significant heart block/bradycardia, or need for preoperative mechanical circulatory/ventilatory support. Inability to complete postoperative interviews, or pregnancy/lactation.

Intervention groups

Patients in the intervention group will receive dexmedetomidine (1 $\mu\text{g}/\text{kg}$ loading dose over 10 minutes, followed by 0.2–0.7 $\mu\text{g}/\text{kg}/\text{hour}$ infusion) along with propofol infusion for maintenance of anesthesia during elective CABG. Propofol will be titrated according to the institutional anesthesia protocol to maintain an adequate depth of anesthesia. Standard intraoperative monitoring and anesthetic management will be the same for both groups.

Main outcome variables

Primary Outcome Variable Intraoperative awareness, assessed using the Modified Brice Interview at 24 and 72 hours after surgery.

General information

Reason for update

Acronym

RCT

IRCT registration information

IRCT registration number: **IRCT20240901062926N2**

Registration date: **2026-07-20, 1405/04/29**

Registration timing: **registered_while_recruiting**

Last update: **2026-07-20, 1405/04/29**

Update count: **0**

Registration date

2026-07-20, 1405/04/29

Registrant information

Name

Muhammad Tayyeb

Name of organization / entity

Superior University

Country

Pakistan

Phone

+92 307 8625941

Email address

tayyabm851@gmail.com

Recruitment status

Recruitment complete

Funding source

Expected recruitment start date

2026-04-22, 1405/02/02

Expected recruitment end date

2026-07-25, 1405/05/03

Actual recruitment start date

empty

Actual recruitment end date

empty

Trial completion date

empty

Scientific title

Comparison of Intraoperative Awareness with Dexmedetomidine-Propofol Infusion versus Intravenous Xylocaine-Propofol Infusion in Patients Undergoing Coronary Artery Bypass Grafting: A Randomized Controlled Trial

Public title

Intraoperative Awareness with Dexmedetomidine-Propofol Infusion versus Intravenous Xylocaine-Propofol Infusion in Patients Undergoing Coronary Artery Bypass Grafting: A Randomized Controlled Trial

Purpose

Prevention

Inclusion/Exclusion criteria**Inclusion criteria:**

Adult patients aged 30–75 years. Patients of either gender. Patients scheduled for elective isolated Coronary Artery Bypass Grafting (CABG) under general anesthesia American Society of Anesthesiologists (ASA) physical status III or IV. Left ventricular ejection fraction (LVEF) $\geq 35\%$ Patients who are able to understand the study protocol and provide written informed consent. Patients who are able to communicate postoperatively and complete the Modified Brice Interview.

Exclusion criteria:

Patients undergoing emergency CABG surgery. Patients scheduled for combined cardiac procedures (e.g., CABG with valve replacement or other cardiac surgeries). History of previous cardiac surgery (redo CABG). Known hypersensitivity or contraindication to dexmedetomidine, propofol, lidocaine (xylocaine), or any study medication. Pre-existing neurological disorders (e.g., stroke with cognitive impairment, epilepsy, dementia) or psychiatric illnesses that may interfere with the assessment of awareness. Chronic use of sedatives, opioids, alcohol, or recreational drugs. Severe hepatic dysfunction (Child-Pugh Class C) or severe renal impairment requiring dialysis. Significant conduction abnormalities (e.g., second- or third-degree atrioventricular block without a permanent pacemaker) or severe bradycardia (heart rate < 50 beats/min). Patients requiring preoperative mechanical ventilation or intra-aortic balloon pump support. Patients with hearing impairment, language barriers, or inability to participate in postoperative interviews. Pregnant or lactating women.

AgeFrom **30 years** old to **75 years** old**Gender**

Both

Phase

0

Groups that have been masked

- Participant

Sample sizeTarget sample size: **100****Randomization (investigator's opinion)**

Randomized

Randomization description

Eligible patients who provide written informed consent will be randomly allocated in a 1:1 ratio to either the Dexmedetomidine Propofol group (Group D) or the Xylocaine-Propofol group (Group X) using a computer-generated randomization sequence. Group allocation will be concealed using sequentially numbered, opaque, sealed envelopes, which will be opened immediately before induction of anesthesia by an anesthesiologist not involved in postoperative outcome assessment. This process will ensure unbiased allocation of participants to the two study groups.

Blinding (investigator's opinion)

Single blinded

Blinding description

This study will be conducted as a single-blinded randomized controlled trial. The participants will be unaware of the treatment group to which they are assigned (Dexmedetomidine-Propofol or Intravenous Xylocaine-Propofol). An anesthesiologist who is not involved in data collection will prepare and administer the study medications according to the randomization sequence. The investigator responsible for postoperative assessment of intraoperative awareness using the Modified Brice Interview and data analysis will remain blinded to the group allocation. The anesthesiologist administering anesthesia cannot be blinded due to the differences in drug preparation and administration protocols. This blinding approach is intended to minimize assessment bias while ensuring patient safety throughout the perioperative period.

Placebo

Not used

Assignment

Parallel

Other design features

none

Secondary Ids

empty

Ethics committees1**Ethics committee****Name of ethics committee**

NATIONAL HOSPITAL & MEDICAL CENTER

Street address

national hospital

City

lahore

Postal code

2246

Approval date

2025-05-25, 1404/03/04

Ethics committee reference number

REF: NHMC/IRB/114

Health conditions studied**1****Description of health condition studied**

Intraoperative awareness is the unintended recall of events experienced during general anesthesia. It is a rare but serious complication that can result in psychological distress, anxiety, sleep disturbances, and post-traumatic stress disorder (PTSD). Patients undergoing coronary artery bypass grafting (CABG) are at a higher risk of intraoperative awareness because anesthetic doses are often reduced to maintain cardiovascular stability. This study aims to compare the effectiveness of dexmedetomidine-propofol infusion versus intravenous xylocaine-propofol infusion in reducing the incidence of intraoperative awareness in patients undergoing elective CABG.

ICD-10 code

Y93

ICD-10 code description

Activity codes

Primary outcomes**1****Description**

Intraoperative awareness, assessed using the Modified Brice Interview conducted at 24 hours and 72 hours after surgery. Patients reporting explicit recall of intraoperative events will be considered to have experienced intraoperative awareness.

Timepoint

24 hour and 72 hours

Method of measurement

Modified Brice Interview Interpretation0 No intraoperative awareness (no explicit recall of intraoperative events).1 Possible intraoperative awareness (uncertain or vague recall requiring further evaluation).2 Definite intraoperative awareness (clear explicit recall of intraoperative events).Interpretation:0: No awareness1: Possible awareness2: Definite awareness

Secondary outcomes

empty

Intervention groups**1****Description**

Intervention group:

Category

Early detection

Recruitment centers**1****Recruitment center****Name of recruitment center**

NATIONAL HOSPITAL & MEDICAL CENTER

Full name of responsible person

Dr. Ahmad Tariq

Street address

national hospital

City

lahore

Postal code

2246

Phone

+92 333 4699734

Fax

+92 333 4699734

Email

drahmed6196@gmail.com

Web page address

<https://nationalhospital.org/>

Sponsors / Funding sources**1****Sponsor****Name of organization / entity**

National Hospital & Medical Center

Full name of responsible person

National Hospital & Medical Center

Street address

national hospital

City

lahore

Postal code

2246

Phone

+92 307 8625941

Fax

+92 307 8625941

Email

tayyabm851@gmail.com

Web page address

<https://nationalhospital.org/>

Grant name

<https://nationalhospital.org/>

Grant code / Reference number**Is the source of funding the same sponsor organization/entity?**

Yes

Title of funding source

National Hospital & Medical Center

Proportion provided by this source

1

Public or private sector

Private

Domestic or foreign origin

Domestic
Category of foreign source of funding
empty
Country of origin
Type of organization providing the funding
Other

Person responsible for general inquiries

Contact

Name of organization / entity
National Hospital & Medical Center
Full name of responsible person
Dr Ahmad Tariq
Position
consultant
Latest degree
Subspecialist
Other areas of specialty/work
Anesthesiology
Street address
national hospital
City
lahore
Province
Punjab
Postal code
2246
Phone
+92 307 8625941
Fax
+92 307 8625941
Email
drahmed6196@gmail.com
Web page address
<https://nationalhospital.org/>

Person responsible for scientific inquiries

Contact

Name of organization / entity
National Hospital & Medical Center
Full name of responsible person
Dr Ahmad Tariq
Position
Consultant
Latest degree
Subspecialist
Other areas of specialty/work
Anesthesiology
Street address
national hospital
City
lahore
Province
Punjab
Postal code
2246
Phone
+92 307 8625941
Fax

+92 307 8625941
Email
drahmed6196@gmail.com
Web page address
<https://nationalhospital.org/>

Person responsible for updating data

Contact

Name of organization / entity
National Hospital & Medical Center
Full name of responsible person
Dr Ahmad Tariq
Position
Consultant
Latest degree
Subspecialist
Other areas of specialty/work
Anesthesiology
Street address
national hospital
City
lahore
Province
Punjab
Postal code
2246
Phone
+92 307 8625941
Fax
+92 307 8625941
Email
drahmed6196@gmail.com
Web page address
<https://nationalhospital.org/>

Sharing plan

Deidentified Individual Participant Data Set (IPD)

Yes - There is a plan to make this available

Study Protocol

Yes - There is a plan to make this available

Statistical Analysis Plan

Yes - There is a plan to make this available

Informed Consent Form

Yes - There is a plan to make this available

Clinical Study Report

Not applicable

Analytic Code

Not applicable

Data Dictionary

Not applicable

Title and more details about the data/document

DATA

When the data will become available and for how long

6 month

To whom data/document is available

open access

Under which criteria data/document could be used

educational purpose

From where data/document is obtainable

National Hospital
**What processes are involved for a request to access
data/document**

email
Comments
thankful